



© CC Somya Pareek, Tejinder Singh Ajmani, Mahima Batra, 2025
<https://doi.org/10.24884/2078-5658-2025-22-4-61-66>

Исследование соотношения окружности шеи и тироментальной дистанции для прогнозирования сложности интубации у пациентов с плановыми операциями в условиях общей анестезии

SOMYA PAREEK*, TEJINDER SINGH AJMANI, MAHIMA BATRA

Отделение анестезии Народного колледжа медицинских наук и исследовательского центра, Бхопал, Мадхья-Прадеш, Индия

Поступила в редакцию 16.05.2025 г.; дата рецензирования 17.06.2025 г.

РЕЗЮМЕ

Введение. Всесторонняя оценка состояния дыхательных путей требует знания анатомических особенностей шеи и верхних дыхательных путей. Точное прогнозирование потенциально трудных дыхательных путей является задачей, зависящей от факторов, связанных с пациентом, клинической обстановкой и квалификацией анестезиолога. Неспособность предвидеть трудности с обеспечением проходимости дыхательных путей остается распространенной причиной осложнений, связанных с анестезией.

Цель – оценить взаимосвязь между окружностью шеи (ОШ), тироментальной дистанцией (ТМД) и частотой возникновения трудностей при интубации у взрослых пациентов, которым выполнены плановые оперативные вмешательства в условиях общей анестезии.

Материалы и методы. В исследование были включены 100 пациентов, относящихся к I и II степени риска по ASA, перенесших плановые операции в условиях общей анестезии с интубацией трахеи. Пациенты были обследованы до операции и во время операции. Сбор данных проводили с использованием анкеты пациента, которая включает демографические данные, индекс массы тела (ИМТ), ТМД, соотношение ОШ/ТМД и модифицированный тест Маллампасти. Для статистического анализа данных использовали статистическое программное обеспечение (SPSS 20.0).

Результаты. В случаях трудной интубации ТМД значительно меньше (среднее = 6,07 см, SD = 0,892), чем при ее отсутствии (7,31 см, SD = 0,877, $P = 0,02994$). Значения соотношения между окружностью шеи и ТМД у лиц без трудной интубации более низкие: 4,78 (SD = 0,466) против 5,71; SD = 0,183 ($P < 0,001$).

Закключение. Соотношение ОШ/ТМД является простым, эффективным и неинвазивным показателем сложности интубации. Оно демонстрирует превосходную чувствительность и специфичность по сравнению с ОШ и Маллампасти-тестом. Регулярное включение этого измерения в предоперационную оценку состояния дыхательных путей может повысить безопасность анестезии.

Ключевые слова: модифицированная шкала Маллампасти, сложная интубация, окружность шеи, тироментальная дистанция, индекс массы тела

Для цитирования: Somya Pareek, Tejinder Singh Ajmani, Mahima Batra Исследование соотношения окружности шеи и тироментальной дистанции для прогнозирования сложности интубации у пациентов с плановыми операциями в условиях общей анестезии // Вестник анестезиологии и реаниматологии. – 2025. – Т. 22, № 4. – С. 61–66. <https://doi.org/10.24884/2078-5658-2025-22-4-61-66>.

A study of the neck circumference to thyromental distance ratio to predict difficult intubation in patients scheduled for elective surgeries under general anaesthesia

SOMYA PAREEK*, TEJINDER SINGH AJMANI, MAHIMA BATRA

Department of Anaesthesia, People's College of Medical Sciences & Research Centre, Bhopal, Madhya Pradesh, India

Received 16.05.2025; review date 17.06.2025

ABSTRACT

Introduction. A comprehensive airway assessment requires an understanding of the anatomical features of the neck and upper airway. Accurately predicting a potentially difficult airway is a multifactorial challenge, dependent on patient-related factors, clinical context, and the anesthesiologist's proficiency. Failure to anticipate a difficult airway remains a common cause of anesthesia-related complications.

The objective was to evaluate the relationship between the neck circumference to thyromental distance ratio (NC/TMD) and the incidence of difficult intubation in adult patients scheduled for elective surgeries under general anesthesia.

Materials and method. This study included 100 patients undergoing elective surgeries under general anesthesia with endotracheal intubation of ASA grades I and II. Patients were examined preoperatively and intraoperatively. Data collection using the patient proforma which includes the demographic data, body mass index (BMI), NC/TMD ratio, thyromental distance (TMD), and Modified Mallampati Test (MMT). Statistical software (SPSS 20.0) was used for statistical analyses of the data.

Results. In cases of difficult intubation, TMD was significantly lower (mean = 6.07 cm, SD = 0.892) than in its absence (7.31 cm, SD = 0.877, $P = 0.02994$). The values of the ratio between neck circumference and TMD in individuals without difficult intubation were lower: 4.78 (SD = 0.466) versus 5.71; SD = 0.183 ($P < 0.001$).

Conclusion. The NC/TMD ratio is a simple, effective, and non-invasive predictor of difficult intubation. It demonstrates superior sensitivity and specificity compared to NC and MMT. Routine incorporation of this measurement in preoperative airway assessment could improve the safety of anesthesia.

Keywords: Modified Mallampati test (MMT), difficult intubation (DI), neck circumference (NC), thyromental distance (TMD), body mass index (BMI)

For citation: Somya Pareek, Tejinder Singh Ajmani, Mahima Batra. A study of the neck circumference to thyromental distance ratio to predict difficult intubation in patients scheduled for elective surgeries under general anaesthesia. *Messenger of Anesthesiology and Resuscitation*, 2025, Vol. 22, № 4, P. 61–66. (In Russ.). <https://doi.org/10.24884/2078-5658-2025-22-4-61-66>.

* Для корреспонденции:
Somya Pareek
E-mail: somyapareek97@gmail.com

* Correspondence:
Somya Pareek
E-mail: somyapareek97@gmail.com

Introduction

The occurrence of a challenging laryngoscopy is approximately the same (around 10%) across non-obese individuals and obese individuals. However, there have been an increasing number of studies indicating challenges in performing intubation in obese patients [10]. This is attributed to difficult visualization of larynx in obese patients which might lead to difficult intubation. DI is a major concern for anesthesiologists and it accounts to 17% of airway injuries and leads to significant perioperative morbidity and mortality. Radiological imaging like ultrasonography, magnetic resonance imaging and computed tomography scans also have shown the presence of excess soft tissue in various areas of neck like the pharynx, retro pharynx, suprasternal notch, at the level of cords and lateral neck region of obese patients [2]. Several clinical indicators can raise the risk of a challenging airway in obese people. Some of these characteristics include an enlarged airway management in patients planned for general anaesthesia is of utmost importance for an anesthesiologist since ages.

Prediction of difficult intubation is one of the most challenging tasks for any anaesthesiologist and it is affected by not only the patient factors but the clinical settings, risk assessment, clinical judgement and skills of an anesthesiologist as well [13]. The difficult endotracheal intubation as per the ASA guidelines is defined as «more than three attempts or more than ten minutes proper for insertion of a tracheal tube with conventional laryngoscopy» [8].

The hypothesis is that the ratio of neck circumference to thyromental distance (NC/TMD ratio) of more than or equal to five will predict a difficult intubation and will have a better statistical and clinical significance as compared to other standard indices of airway assessment among patients. Therefore, in this study, our goal was to investigate a preoperative indicator of challenging intubation named the ratio of neck circumference to thyromental distance.

This indicator does not require any specialized equipment, can be performed quickly, is not uncomfortable for the patient, and is noninvasive. The current study examines the efficacy of the neck circumference/thyromental distance ratio (NC/TM distance ratio) in comparison to the Mallampati score and neck circumference as accurate indicators for predicting difficult intubation.

The objective of the study was to assess the validity of neck circumference to thyromental distance ratio (NC/TMD) as an important parameter to predict difficult intubation amongst adult patients undergoing elective surgery under general anaesthesia.

Materials and methods

The clinical study was conducted at People's College of Medical Sciences & Research Centre, Bhopal, India in the span of six months.

Permission of the Ethics Committee: this work was approved by Institutional Ethical Committee Code: IEC-2022/66.

100 patients were studied after approval by the Institutional Ethical Committee (IEC-2022/66) and written informed consent. Age more than 18 and less than 65 years, ASA class I and II, patients scheduled for elective surgery under general anaesthesia with endotracheal intubation were included.

Exclusion criteria: patients less than 18 years and more than 65 years of age. Patients scheduled for general anaesthesia without endotracheal intubation such as mask ventilation, use of laryngeal mask airways, sedation, monitored anesthetic care were excluded. Patients coming for surgery under regional anaesthesia. Patients with upper airway pathologies, thyroid gland dysfunction, facial and maxillary fractures. Obstetric patients and cervical spine fractures were excluded along with patient refusal to participate in the study.

Preoperative assessment. It consisted of proforma with emphasis on a) demography of patient b) body mass index c) airway assessment variables include: 1) neck circumference (measured at the level of cricoid cartilage); 2) thyromental distance (the distance between thyroid cartilage and mentum with neck fully extended); 3) the ratio of neck circumference to thyromental distance (NC/TM distance ratio); 4) Mallampati classification without phonation.

Intraoperative assessment. Difficulty of intubation was assessed by anesthetist by filing up intubation difficulty score (validated IDS score) [17 Adnet] after intubation. Intubation difficulty score consists of seven variables from N1 to N7. The sum of N1 to N7 gives total IDS score. Any score of greater than or equal to five was considered to be difficult intubation and score less than five considered to be easy intubation.

Statistical analysis. The results are presented as numbers (percentage). The Chi-square test was used. Statistical analysis was performed using SPSS software, and statistical significance was set at $p < 0.05$.

Results

The distribution illustrates a clear trend in table 1: as the Mallampati grading increases, the proportion of difficult intubation cases also rises, particularly noticeable in Grades 3 and 4, where the rates significantly exceed those of Grades 1 and 2. The statistical significance of this trend was evaluated using Pearson's Chi-square test, resulting in a Chi2 value of 26.67 and a P-value of less than 0.0001. This highly significant P-value indicates a strong association between higher Mallampati grades and an increased incidence of difficult intubation.

Table 2 categorizes the ease of intubation according to the Intubation Difficulty Scale (IDS) and its correlation with the occurrence of difficult intubation among 100 participants.

Table 3 provides an analysis of various body parameters and their correlation with difficult intubation among participants, summarized as follows:

Table 1. Mallampati Grading among participants (*n* = 100)

MPC	Difficult Intubation		
	NO	YES	Total
1	33	3	36
	40.24	16.67	36.00
2	41	4	45
	50.00	22.22	45.00
3	6	6	12
	7.32	33.33	12.00
4	2	5	7
	2.44	27.78	7.00
Total	82	18	100

Pearson Chi2 = 26.67; P-value < 0.0001




Fig 1. Mallampati Grading among participants (*n* = 100)




Fig 2. Ease of Intubation according to IDS Scale (*n* = 100)

Table 2. Ease of Intubation according to IDS Scale (*n* = 100)

IDS	Difficult Intubation		
	No	YES	Total
Easy	18	0	18
	21.95	0.00	18.00
Slightly Difficult	64	10	74
	78.05	55.56	74.00
Moderate to Major Difficult	0	8	8
	0.00	44.44	8.00
Total	82	18	100

Pearson Chi2 = 41.40; *p*-value < 0.0001

- BMI (Body Mass Index): the mean BMI for participants without difficult intubation was 25.2 (SD = 2.67), while for those with difficult intubation, it was 28.00 (SD = 2.40). The difference is statistically significant (P-value = 0.0001), indicating higher BMI is associated with an increased incidence of difficult intubation;

- Obesity: 7.2% of participants without difficult intubation were classified as obese, compared to 22.2% of those with difficult intubation. The difference approached statistical significance (P-value = 0.059), suggesting a trend toward higher obesity rates in those experiencing difficult intubation;

- Neck circumference: The mean neck circumference for participants without difficult intubation was 34.96

cm (SD = 5.44), and for those with difficult intubation, it was significantly higher at 40.46 cm (SD = 5.68) (P-value = 0.0002).

Thyromental distance: there was a significant difference in thyromental distance between those without (mean = 7.31 cm, SD = 0.877) and with difficult intubation (mean = 6.07 cm, SD = 0.892) (P-value = 0.02994). Neck Circumference-to-Thyromental Distance Ratio: participants without difficult intubation had a lower mean ratio of 4.78 (SD = 0.466) compared to those with difficult intubation, who had a significantly higher mean ratio of 5.71 (SD = 0.183) (P-value < 0.001). The 95% confidence intervals for these means did not overlap, further highlighting the significant difference.

Table 3. Body Parameters

Parameter Mean (SD)	Difficult Intubation		P-value
	No	YES	
BMI	25.2 (2.67)	28.00 (2.40)	0.0001
Obese	6 (7.2%)	4 (22.2%)	0.059
NC	34.96 (5.44)	40.46(5.68)	0.0002
TMD	7.31 (0.877)	6.07 (0.892)	0.029
NC/TMD	4.78 (0.466)	5.71 (0.183)	< 0.0001

Table 4. Logistic regression for difficult intubation

Variable	Odds ratio	P-value	95% CI	
NC/TMD	9.47	< 0.0001	3.30	25.86
BMI	0.85	0.426	0.58	1.25
Age	0.963	0.324	0.89	1.03
Male	0.909	0.917	0.15	5.49
Female	1.0	–	–	–

The analysis indicates that certain body parameters, particularly BMI, neck circumference, and the neck circumference-to-thyromental distance ratio, are significantly associated with an increased risk of difficult intubation. These findings underscore the importance of considering these parameters in the preoperative assessment to identify patients at risk for difficult intubation and prepare appropriate airway management strategies.

Table 4 presents the results of a logistic regression analysis investigating the association of various factors with the likelihood of difficult intubation, expressed as odds ratios (ORs), P-values, and 95% confidence intervals (CIs).

Neck Circumference-to-Thyromental Distance Ratio: the analysis showed a significant association, with an OR of 9.47 (95% CI: 3.30–25.86, P-value < 0.0001), indicating that each unit increase in the ratio increases the odds of difficult intubation by approximately 9.47 times.

- BMI (Body Mass Index): BMI was not significantly associated with difficult intubation, with an OR of 0.85 (95% CI: 0.58–1.25, P-value = 0.426). This suggests that, within this study, BMI alone does not significantly influence the likelihood of difficult intubation.

- Age: age also showed no significant association with difficult intubation, evidenced by an OR of 0.963 (95% CI: 0.89–1.03, P-value = 0.324), indicating that the likelihood of difficult intubation does not significantly decrease or increase with age in this sample.

- Gender: male participants had an OR of 0.909 (95% CI: 0.15–5.49, P-value = 0.917) compared to female participants, indicating no significant gender difference in the odds of experiencing difficult intubation. Female participants are the reference category (OR = 1.0). The results from the logistic regression analysis underscore the importance of the neck circumference-to-thyromental distance ratio as a significant predictor of difficult intubation. In contrast, BMI, age, and gender were not statistically significant predictors in this model. This analysis provides valuable insights into factors that may warrant consideration during preoperative

airway assessments to identify patients at higher risk for difficult intubation.

Discussion

The existing data are inconclusive and there is a divergence of perspectives on the challenge of tracheal intubation. Various endeavors have been made to provide precise predictors for difficulty laryngoscopy or difficult intubation. The present study was undertaken to assess the validity of neck circumference to thyromental distance ratio (NC/TMD) as an important parameter to predict difficult intubation amongst adult patients undergoing elective surgery under general anaesthesia.

A total of 100 patients who were scheduled for elective surgery were evaluated for the presence of difficult intubation in the study. The group consisted of individuals from varied demographic backgrounds and with varying physical attributes, enabling a thorough investigation of factors that predict and the occurrence rate of difficult intubation. Patients had a preoperative assessment, which included the evaluation of the Mallampati score, measurement of thyromental distance, and assessment of neck mobility. Anesthesia was delivered according to established protocols, and the level of difficulty in intubation was assessed using the Intubation Difficulty Scale (IDS). The findings indicated that 18% of patients encountered challenging intubation, specifically defined as having an IDS score of 5 or higher. The reported incidence by G.Fotopoulou et al. [7] (20%), G.S. Voyagis et al. [24] (20.2%), D. Castro et al. [4] (20.75%). In contrast, other authors such as T. Shiga et al. [21] (15.8%), P.Juvin et al. [12] (15.5%), H.Gonzalez et al. [9] (14.3%), W.H. Kim et al. [13] (13.8%), and S.Shailaja et al. [22] (11%) reported a lower incidence. Therefore, the literature study indicates that the occurrence of problematic intubation ranged from 11% to 22%.

The present study found a highly significant P-value ($p < 0.0001$) indicating a strong association between higher Mallampati grades and an increased incidence of difficult intubation. The distribution of patients ac-

cording Mallampati grade score illustrates a clear trend: as the Mallampati grading increases, the proportion of difficult intubation cases also rises, particularly noticeable in Grades 3 and 4, where the rates significantly exceed those of Grades 1 and 2. Our results are in concordance with other studies reported in the literature. The Mallampati score has become a standard part of a comprehensive airway evaluation, although its predictive value for difficult intubation has proven to be low. The Mallampati score is based upon visualization of anatomical oropharyngeal structures and relates them to intubation difficulty [20]. One of the adjustments made to the Mallampati scoring system is the one by Samsoon and Young, which adds a fourth class to the descriptions of oropharyngeal exposures [20]. Some drawbacks of the modified Mallampati classification include interobserver variability and reliance on patient compliance. According to research by A. Lee et al. [14], the categorization was not very good at predicting difficult intubations. Additionally, other authors have found that, in order to improve diagnostic accuracy, it should be used in conjunction with other predictors rather than being used alone as a test to predict difficult intubation or laryngoscopy [16, 18].

The Intubation Difficulty Scale (IDS) was developed by F. Adnet et al. [1]. It has seven objective features that are utilized for the assessment of intubation. IDS has been employed as a verified measure of difficulty to characterize challenging intubation. The present study utilized the F. Adnet Intubation Difficulty Scale score to determine the frequency of intubation difficulty in individuals undergoing elective surgery under general anesthesia.

W. Koh et al. [14] found the Cormack-Lehane grade was most sensitive for predicting difficult intubation compared with other parameters of IDS and their findings indicated a correlation between Cormack-Lehane grade 3 or 4 and the likelihood of encountering challenges during intubation. These findings suggest that even though the actual incidence of difficult intubation (5.8%) was lower in patients with a Cormack-Lehane grade of 3 or 4 (16.2%), anesthesiologists should always be prepared for difficult intubation when encountering patients with a high Cormack-Lehane grade.

The present study found a clear delineation between the perceived difficulty of intubation and the actual incidence of difficult intubation, especially highlighted by the 100% incidence rate in the «Moderate to Major Difficult» category with a statistical significance ($p = 0.0000$) difference. This highly significant P-value strongly indicates a significant association between the IDS category and the occurrence of difficult intubation. All 18 cases classified as «Easy» experienced no difficult intubation, indicating a 0% incidence of diffi-

culty in this category. Of the 74 cases deemed «Slightly Difficult», 10 were identified as difficult intubations, accounting for 55.56% of all difficult intubation cases. In the «Moderate to Major Difficult» category, all 8 cases were considered difficult intubations, comprising 44.44% of the difficult cases. The present study analyzed various body parameters and their correlation with difficult intubation among participant. The mean BMI for participants without difficult intubation was 25.2 (SD = 2.67), while for those with difficult intubation, it was 28.00 (SD = 2.40). The difference is statistically significant (P-value = 0.0001), indicating higher BMI is associated with an increased incidence of difficult intubation. The mean neck circumference for participants without difficult intubation was 34.96 cm (SD = 5.44), and for those with difficult intubation, it was significantly higher at 40.46 cm (SD = 5.68) (P-value = 0.0002). There was a significant difference in thyromental distance between those without (mean = 7.31 cm, SD = 0.877) and with difficult intubation (mean = 6.07 cm, SD = 0.892) (P-value = 0.02994). Participants without difficult intubation had a lower mean ratio of 4.78 (SD = 0.466) compared to those with difficult intubation, who had a significantly higher mean ratio of 5.71 (SD = 0.183) (P-value < 0.001). The 95% confidence intervals for these means did not overlap, further highlighting the significant difference. Comparing our study with various studies, a study conducted by W. Ittichaikulthol et al. [11] with a substantial sample size that the combination of MMT (Mallampati Modified Test) and TMD (Thyromental Distance) were reliable indicators of a challenging laryngoscopy procedure in the Thai population. Nevertheless, they employed a TMD value of < 6 cm which differs from the < 6.5 cms parameter utilized in the present investigation [11]. Another study by E. Magalhães et al. [17] that evaluated Chinese women who were pregnant and those who were not pregnant and discovered that the combination of predicting factors might enhance the accuracy. A study conducted by J. L. Díaz-Gómez et al. in the United States found that the utilization of up to ten measurements significantly enhanced the predictability of a challenging intubation by improving the aggregate set of factors [6].

The results from the logistic regression analysis underscore the importance of the neck circumference-to-thyromental distance ratio as a significant predictor of difficult intubation. In contrast, BMI, age, and gender were not statistically significant predictors in this model. This analysis provides valuable insights into factors that may warrant consideration during preoperative airway assessments to identify patients at higher risk for difficult intubation.

Conflict of Interest. The authors declare no conflicts of interest.

Contribution of the authors. All authors made a substantial contribution to the conception of the work, acquisition, analysis, interpretation of data for the work, drafting and revising the work, and final approval of the version to be published, and agreed to be accountable for all aspects of the work.

REFERENCES

- Adnet F, Borron S, Racine S. The intubation difficulty scale (IDS): proposal and evaluation of a new score characterizing the complexity of endotracheal intubation. *Anaesthesiology*, 1997, 87, pp. 1290–1297. <http://doi.org/10.1097/0000542-199712000-00005>.
- Ankalwar V. R., Patel M., Tirpude N. G. Neck circumference to thyromental distance ratio: Is a reliable predictor of difficult intubation in obese patients. *Indian J Clin Anaesth*, 2019, vol. 6, no. 1. <http://doi.org/152-6.10.18231/2394-4994.2019.0029>.
- Benumof J. L. Management of difficult adult airway with special emphasis on awake tracheal intubation. *Anesthesiology*, 1991, vol. 75, pp. 1087–1110. <http://doi.org/10.1097/0000542-199112000-00021>.
- Castro D., Leão P., Pacheco M. et al. Neck circumference to thyromental distance ratio: evaluation of a new predictive tool of difficult intubation in obese patients submitted to bariatric surgery: 19AP6-8. *European Journal of Anaesthesiology*, 2013, vol. 30, pp. 268–269.
- Chara L., Eleftherios V., Maria M. et al. Anatomic features of the neck as predictive markers of difficult direct laryngoscopy in men and women: A prospective study. *Indian journal of anaesthesia*, 2014, vol. 58, no. 2, pp. 176–182. <http://doi.org/10.4103/0019-5049.130822>.
- Díaz-Gómez J. L., Satyapriya A., Satyapriya S. V. et al. Standard clinical risk factors for difficult laryngoscopy are not independent predictors of intubation success with the GlideScope. *J Clin Anesth*, 2011, vol. 23, pp. 603–610. <http://doi.org/10.1016/j.jclinane.2011.03.006>(94).
- Fotopoulou G., Vasileiou I., Dre K. et al. Can we predict difficult intubation in obese patients: 19AP4-7. *European Journal of Anaesthesiology*, 2011, vol. 28, pp. 234. <http://doi.org/10.47009/jamp.2024.6.6.127>.
- Frerk C., Mitchell V. S., McNarry A. F. et al. Difficult Airway Society 2015 guidelines for management of unanticipated difficult intubation in adults. *British Journal of Anaesthesia*, 2015, vol. 115, no. 6, pp. 827–848. <http://doi.org/10.1093/bja/aev371>.
- Gonzalez H., Minville V., Delanoue K. et al. The importance of increased neck circumference to intubation difficulties in obese patients. *Anesth Analg*, 2008, vol. 106, no. 4, pp. 1132–1136. <http://doi.org/10.1213/ane.0b013e3181679659>.
- Hirmanpour A., Safavi M., Honarmand A. et al. The predictive value of the ratio of neck circumference to thyromental distance in comparison with four predictive tests for difficult laryngoscopy in obstetric patients scheduled for caesarean delivery. *Advanced biomedical research*, 2014, vol. 3, no. 1, pp. 2000. <http://doi.org/10.4103/2277-9175.142045>.
- Ittichakulthol W., Chanpradub S., Amnondetchakorn S. et al. Modified Mallampati test and thyromental distance as a predictor of difficult laryngoscopy in Thai patients. *J Med Assoc Thai*, 2010, vol. 93, pp. 84–89. PMID: 20196416.
- Juvin P., Lavaut E., Dupont H. et al. Difficult tracheal intubation is more common in obese than in lean patients. *Anesth Analg*, 2003, vol. 97, pp. 595–600. <http://doi.org/10.1213/01.ANE.0000072547.75928.B0>.
- Kim W. H., Ahn H. J., Lee C. J. et al. Neck circumference to thyromental distance ratio: a new predictor of difficult intubation in obese patients. *British journal of anaesthesia*, 2011, vol. 106, no. 5, pp. 743–748. <http://doi.org/10.1093/bja/aer024>.
- Koh W., Kim H., Kim K. et al. Encountering unexpected difficult airway: relationship with the intubation difficulty scale. *Korean journal of anesthesiology*, 2016, vol. 69, no. 3, pp. 244–9. <http://doi.org/10.4097/kjae.2016.69.3.244>.
- Lee A., Fan L. T., Gin T. et al. A systematic review (meta analysis) of the accuracy of the mallampatti tests to predict the difficult airway. *Anesth Analg*, 2006, vol. 102, pp. 1867–1878. <http://doi.org/10.1213/01.ane.0000217211.12232.55>.
- Lundström L. H. Detection of risk factors for difficult tracheal intubation. *Dan Med J*, 2012, vol. 59, B4431. <http://doi.org/10.2174/1874321801610010034>(86).
- Magalhães E., Oliveira M. F., Sousa G. C. et al. Use of simple clinical predictors on preoperative diagnosis of difficult endotracheal intubation in obese patients. *Braz J Anesthesiol*, 2013, vol. 63, no. 3, pp. 262–266. [http://doi.org/10.1016/S0034-7094\(13\)70228-9](http://doi.org/10.1016/S0034-7094(13)70228-9).
- Rose N., Maya G., Sathiyadas A. Comparison of difficult intubation and neck circumference to thyromental distance ratio, in obese and non-obese: A clinical study. *J Med Sci Clin Res*, 2017, vol. 5, pp. 19670–19679. <http://doi.org/10.2174/1874321801610010034>.
- Rosenberg M. B., Phero J. C. Airway assessment for office sedation/anesthesia. *Anesth Prog*, 2015, vol. 62, no. 2, pp. 74–80. <http://doi.org/10.2344/0003-3006-62.2.74>.
- Samsoon G. L., Young J. R. Difficult tracheal intubation: a retrospective study. *Anaesthesia*, 1987, vol. 42, no. 5, pp. 487–90. [http://doi.org/10.1111/j.1365-2044.1987.tb04039.\(77\)](http://doi.org/10.1111/j.1365-2044.1987.tb04039.(77)).
- Shiga T., Wajima Z., Inoue T., Sakamoto A. Predicting difficult intubation in apparently normal patients: A meta-analysis of bedside screening test performance. *Anesthesiology*, 2005, vol. 103, no. 42, pp. 9–37. <http://doi.org/10.1097/0000542-200508000-00027>.
- Shailaja S., Nichelle S. M., Shetty A. K. et al. Comparing ease of intubation in obese and lean patients using intubation difficulty scale. *Anesthesia Essays and Researches*, 2014, vol. 8, no. 2, pp. 168–174. <http://doi.org/10.4103/0259-1162.134493>.
- Shamim A., Swami S., Konnur S. et al. The importance of evaluation of neck circumference to thyromental distance ratio as a predictor of difficult intubation. *Archives of Anesthesiology and Critical Care*, 2024. <https://doi.org/10.18502/aacc.v10i2.15056>.
- Voyagis G. S., Kyriakis K. P., Dimitriou V. et al. Value of oropharyngeal Mallampati classification in predicting difficult laryngoscopy among obese patients. *Eur J Anaesthesiol*, 1998, vol. 15, pp. 330–334. <http://doi.org/10.1046/j.1365-2346.1998.00301.x>.

INFORMATION ABOUT AUTHORS:

People's University; Department of Anaesthesia, People's College of Medical Sciences & Research Centre, Bhopal, Madhya Pradesh, India

Dr. Somya Pareek

Final Year Resident, Corresponding Author, People's College of Medical Sciences & Research Centre, Bhopal, MP, India.
ORCID: 0009-0007-5891-1631

Dr. Tejinder Singh Ajmani

Associate Professor, Department of Anaesthesia, People's College of Medical Sciences & Research Centre, Bhopal, MP, India.
ORCID: 0009-0001-0656-1912

Dr. Mahima Batra

Head of the department, Professor, Department of Anaesthesia, People's College of Medical Sciences & Research Centre, Bhopal, MP, India.
ORCID: 0000-0002-9734-2810